

098 Weekly TC療法(PTX + CBDCA)

病名: 卵巣がん

1コース: 4週間(3投1休)

申請日 令和2年6月8日
委員会承認日 令和2年7月15日

投与スケジュール

投与方法・投与量			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
末梢メイン																														
①	生理食塩液	100mL 投与時間: 5分	↓							↓							↓													
②	生理食塩液	50mL	↓							↓							↓													
	グラニセトロン3mg	1A	↓							↓							↓													
	ファモチジン20mg	1A	↓							↓							↓													
	デキサート3.3mg【4mg】	2A	↓							↓							↓													
	クロルフェニラミンマレイン酸塩注5m _g	1A	↓							↓							↓													
	投与速度: 200mL/h																													
③	5%ブドウ糖	250mL	↓							↓							↓													
	パクリタキセル	【80mg/m ² 】	↓							↓							↓													
	投与時間: 1時間																													
④	5%ブドウ糖	250mL	↓							↓							↓													
	カルボプラチン	【AUC 2】	↓							↓							↓													
	投与時間: 1時間																													

・CBDCAの投与量(mg/body) = AUC × (GER+25)

GFR = (140 - 年齢) × 体重 / (72 × 血清クレアチニンmg/dl)【女性の場合は: 0.85 ×】

・CBDCAの投与量(mg/body) = AUC × 実側Ccr

・CBDCAの投与量(mg/body) = AUC × 実側Ccr × BSA/1.48