

099 dose-dence weekly TC療法 (PTX+CBDCA)

病名: 卵巣がん

1コース: 4週間(3投1休)

申請日 令和2年6月8日  
委員会承認日 令和2年7月15日

投与スケジュール

| 投与方法・投与量 |                      |                        | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
|----------|----------------------|------------------------|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| 末梢メイン    |                      |                        |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ①        | 生理食塩液                | 100mL                  | ↓ |   |   |   |   |   |   | ↓ |   |    |    |    |    |    | ↓  |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | 投与時間: 5分             |                        |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ②        | 生理食塩液                | 50mL                   | ↓ |   |   |   |   |   |   | ↓ |   |    |    |    |    |    | ↓  |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | グラニセトロン3mg           | 1A                     | ↓ |   |   |   |   |   |   | ↓ |   |    |    |    |    |    | ↓  |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | ファモチジン20mg           | 1A                     | ↓ |   |   |   |   |   |   | ↓ |   |    |    |    |    |    | ↓  |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | デキサート3.3mg【4mg】      | 2A                     | ↓ |   |   |   |   |   |   | ↓ |   |    |    |    |    |    | ↓  |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | ケルフェニミンマレイン酸塩注5mg 1A |                        | ↓ |   |   |   |   |   |   | ↓ |   |    |    |    |    |    | ↓  |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | 投与速度: 200mL/h        |                        |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ③        | 5%ブドウ糖               | 250mL                  | ↓ |   |   |   |   |   |   | ↓ |   |    |    |    |    |    | ↓  |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | パクリタキセル              | 【80mg/m <sup>2</sup> 】 | ↓ |   |   |   |   |   |   | ↓ |   |    |    |    |    |    | ↓  |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | 投与時間: 1時間            |                        |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ④        | 5%ブドウ糖               | 250mL                  | ↓ |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | カルボプラチン              | 【AUC 6】                | ↓ |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
|          | 投与時間: 1時間            |                        |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |

•CBDCAの投与量(mg/body) = AUC × (GER+25)

GFR = (140 - 年齢) × 体重 / (72 × 血清クレアチニンmg/dl)【女性の場合は:0.85 ×】

•CBDCAの投与量(mg/body) = AUC × 実側Ccr

•CBDCAの投与量(mg/body) = AUC × 実側Ccr × BSA/1.48